

APPLICATIONS OF TAYLOR SERIES IN REAL ANALYSIS

Afnan Ali Hasan Alrabiaa^{1*}

Abstract: This paper explores the versatile applications of Taylor series, a powerful mathematical tool employed for approximating complex functions and solving differential equations. While polynomial functions offer simplicity, Taylor series enable the representation of intricate functions, facilitating the study of their asymptotic behavior. Through various examples, this paper demonstrates the diverse applications of Taylor series in approximation and computational science, particularly benefiting engineering and physical science fields. Key advantages include the ability to express any mathematical function as a polynomial, providing approximate solutions in cases where precise solutions are challenging to obtain. In numerical mathematics, Taylor series forms the basis for many equation-solving algorithms, emphasizing its significance. The paper also discusses the importance of considering precision requirements in practical applications. For instance, in scenarios like automatic aircraft landing systems, an error tolerance of one or two meters may be acceptable, while highly precise measurements, such as head position readings from a cylinder, demand errors within a millionth of a meter.

Introduction

Fluid dynamics involves physics principles that concern the behavior of moving fluids, either in consumable or gaseous form and analyses the forces relative to the fluids. The study and analysis of any fluid property and behavior has to employ some rigorous mathematics, and in this case, the Taylor series expansion is very essential. Taylor's series allows the approximation of functions by breaking them down into an infinite number of terms reliant on their derivatives at a specific interval. At a very basic level, this approach aids in fluid property analysis on velocity, pressure, and density by looking at their behavior at a particular point [1].

In fluid dynamics, Taylor series can be used easily and because of its ease, approximation for different functions becomes possible especially if differentiation or integration cannot be done analytically. The conduct of cutting off the series to a given finite number of terms yields better understanding of the fluid flow by forming polynomial approximation to the function. This approach is also very useful in fluid motion governing differential equations, making possible the numerical simulation and stability analysis [2].

In addition, Taylor series together with the concept of fluid parcel's deformation is one of the most important aspects in any flow analysis. In fluid flow analysis, a small change in any parameter can be expressed in its Taylor series to measure how it alters the system. This is very important in objectives such as aerodynamics, meteorology, or hydrodynamics where modeling fluid is critical in accurate forecasting [3].

¹ Assistant Professor, Shaqra University, Common First Year Deanship, Saudi Arabia.
Email: afnanali202020@gmail.com

*Correspondence: afnanali202020@gmail.com

The application of the Taylor series is not only bound to fluid dynamics, but it is also relevant in other branches of science like physics, engineering, or applied mathematics. With the knowledge of the principle of Taylor series expansion, scientists can construct such numerical solutions to differential equations, analyze functions convergence, and study systems' asymptotic behavior. This study focuses on how these series work in fluid dynamics and provides very detailed information on the concept, its usefulness, and importance in both computational and experimental work [4].

One of the most fundamental concepts in fluid dynamics is the expansion of nearby points of vectors and functions. This method can help us understand the properties of fluids. For instance, by examining the nearby parcels' properties (such as density, velocity, pressure, etc.), we can determine their deformations. To gain a deeper understanding of the various features of a fluid, we need to first learn how to expand a function. In the following analysis, we will look closely at the mathematical results. Assume a *power series* representation of a function converges about the point x_o . The form of a power series for a function of x is [16]:

$$f(x) = \sum_{n=0}^{\infty} c_n (x - x_o)^n = c_0 + c_1(x - x_o) + c_2(x - x_o)^2 + c_3(x - x_o)^3 + O((x - x_o)^4)$$

(In the above equation, the expression $O((x - x_o)^4)$ means terms of 4th order and higher.)

This expansion lets in for the approximation of features the use of a finite range of terms, which can be particularly beneficial when dealing with capabilities that are not effortlessly incorporated or differentiated. For example, in numerical evaluation, the Taylor collection is hired to broaden numerical methods for solving differential equations and for performing numerical integration.

One of the full-size applications of the Taylor collection in real evaluation is within the realm of feature approximation. By truncating the Taylor collection after a finite wide variety of phrases, we reap a polynomial that approximates the characteristic close to the factor of enlargement. This polynomial, known as a Taylor polynomial, gives a less difficult feature that carefully resembles the unique characteristic inside a positive c language. The accuracy of this approximation improves as more phrases are protected inside the collection, making it a versatile tool for diverse analytical functions [5].

Another critical software is in the subject of differential equations. Many differential equations that model physical phenomena cannot be solved exactly. The Taylor collection gives a technique to find approximate answers by using increasing the unknown feature as a chain after which solving for the coefficients term via time period. This approach is foundational in perturbation idea, in which small deviations from recognized answers are studied using collection expansions [7].

Furthermore, the Taylor series is instrumental in the take a look at of the convergence houses of capabilities. By reading the rest of the Taylor series, one can determine the radius of convergence and understand the conduct of capabilities within this c program language period. This evaluation is essential in complex evaluation and within the have a look at of analytic features, wherein the capability to specific a function as a power series within a certain domain gives deep insights into its homes [11].

In addition to theoretical programs, the Taylor collection has practical uses in fields consisting of physics and engineering. For example, in physics, the Taylor series is used to approximate the ability power in a device around equilibrium factors, leading to linear approximations that simplify the analysis of physical structures. In

engineering, it aids inside the design and analysis of manage structures, where small deviations from a hard and fast point are studied using linear approximations derived from the Taylor series [19].

In summary, the Taylor collection is a fundamental tool in actual analysis, offering effective strategies for approximating features, solving differential equations, and analyzing the convergence properties of capabilities. Its versatility and practicality make it an critical component of mathematical evaluation and its programs throughout various scientific and engineering disciplines.

The expression means the terms of higher and fourth order. The above extension should converge at the middle value. It means that the sum of the infinite terms must be a finite value. This means that the higher order periods must be reduced in order for this to happen [20]. The coefficients can be examined by comparing the derivatives of the given equation. To define the above extension, it must converge around the center value, x_o . This means that the sum of all infinite terms in the equation above must be a finite value. Therefore, higher order periods must be reduced to make this possible. The coefficients $c_0, c_1, c_2, c_3, \dots, c_n$ can be valued by examining the derivatives of $f(x)$ at the point $x = x_o$.

$$\begin{aligned} f(x)_{x=x_o} &= c_0 \\ \frac{d}{dx} f(x)_{x=x_o} &= f'(x)_{x=x_o} = c_1 \\ \frac{d^2}{dx^2} f(x)_{x=x_o} &= f''(x)_{x=x_o} = 2c_2 \Rightarrow c_2 = \frac{f''(x_o)}{2} \\ f'''(x)_{x=x_o} &= 3 \cdot 2 \cdot c_3 \Rightarrow c_3 = \frac{f'''(x_o)}{3!} \\ f^{(n)}(x)_{x=x_o} &= (n \cdot (n-1) \cdot (n-2) \dots 1) c_n = (n!) c_n \Rightarrow c_n = \frac{f^{(n)}(x_o)}{n!} \end{aligned}$$

The resulting expansion of $f(x)$, with the coefficients derived above, is called the *Taylor series generated by $f(x)$ at $x = x_o$* . Mathematically it is expressed as:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_o)}{n!} (x - x_o)^n = f(x_o) + f'(x_o)(x - x_o) + \frac{f''(x_o)}{2} (x - x_o)^2 + O(x - x_o)^3$$

The selection of a point is not restricted by the original requirement to converge. However, there are practical limitations that prevent us from having many terms to consider when determining the correct equation. For instance, we can only choose a point near the center [17].

To reiterate this point, let us restrict our choice of x_o and the domain of x such that $x - x_o = \delta x < 1$. Then we can express $x = x_o + \delta x$ in equation to obtain

$$f(x_o + \delta x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_o)}{n!} (\delta x)^n = f(x_o) + f'(x_o) \delta x + O(\delta x^2)$$

Since we have chosen δx to be arbitrarily small compared to x_o , it is safe assume the Taylor series approximation is accurate out to the first two terms of the expansion [5, 14].

$$f(x_o + \delta x) \approx f(x_o) + f'(x_o) \delta x$$

The extension of previous equations to multi-variable functions can be tedious. But, if we consider the expansion of only one element at a time, the procedure is simple. We can look at three different cases of previous equation if we consider the variations of only one variable in each set of conditions. The function can be expanded near the point of the three spatial coordinates [21, 22].

$$\begin{aligned}
f(x_o + \delta x, y_o, z_o) &= f(x_o, y_o, z_o) + \left. \frac{\partial f}{\partial x} \right|_{x_o, y_o, z_o} \delta x + O(\delta x^2) \\
&= f(x_o, y_o, z_o) + \left. \frac{\partial f}{\partial y} \right|_{x_o, y_o, z_o} \delta y + O(\delta y^2) \\
&= f(x_o, y_o, z_o) + \left. \frac{\partial f}{\partial z} \right|_{x_o, y_o, z_o} \delta z + O(\delta z^2)
\end{aligned}$$

We can get the vector expression if we consider the variations in the three directions at the same time.

$$\begin{aligned}
f(\vec{x}_o + \delta \vec{x}) &= f(\vec{x}_o) + \nabla f(\vec{x}_o) \cdot \delta \vec{x} + O(|\delta x|^2) \\
\text{where } \vec{x}_o &= x_o \hat{i} + y_o \hat{j} + z_o \hat{k} \text{ and } \delta \vec{x} = \delta x \hat{i} + \delta y \hat{j} + \delta z \hat{k}.
\end{aligned}$$

For most fluid dynamics calculations, the distance from the center of the expansion is assumed to be small enough that the Taylor series can expand out to two terms [10].

Literature Review

The exploration of Taylor series applications in this paper aligns with and expands upon existing research in various scientific domains. While the paper delves into diverse applications of Taylor series, it's crucial to acknowledge the foundational contributions and context provided by previous studies.

Numerical Methods and Approximation

Many researchers have extensively explored the application of Taylor series in numerical methods and function approximation. Works by Veigend, et al. [20] and Bawazeer, et al. [3] provide insights into Taylor series-based numerical integration and radial basis function expansion. The current paper builds upon this foundation by showcasing practical examples in physics, chemistry, and engineering, illustrating the versatility of Taylor series in solving real-world problems.

Numerical methods are very important, especially in mathematical problems that are too difficult to solve and do not have a direct solution. One of the approaches that is popular in useful numerical methods is the Taylor series, which is used in function and integral as well as differential equation estimation. The series expansion permits depiction of a function as a sum of an infinite series of its derivatives, thereby allowing a function to be accurately approximated for many areas of science and engineering.

The use of Taylor series for numerical methods has been the subject of a number of investigations. Amarioarei, et al. [1] for example, proposed efficient nonlinear parsimonious forest models by quicker site productivity and taper analysis through Taylor series expansion. Resources such as Bawazeer, et al. [3] has suggested a new style using radial basis functions based on the partition of unity with Taylor series expansion which has shown to be much more efficient.

Elsaid and Helal [4] in the area of differential equations presented moving Taylor series method for solving one dimensional Stefan problems, which proved to be efficient for phase-change problems. Moreover, Estévez Schwarz and Lamour [5] have used explicit and implicit Taylor series methods for differential-algebraic equations, thus applying the reconstruction Taylor series method in ordinary equations demonstrated the wide ranging possibilities of this approach.

These contributions illustrate the importance of the Taylor series in numerical estimation as an effective and precise technique of working on intricate

mathematical computations across many fields of science.

Definite Integrals and Function Evaluation

Guillot, et al. [7] and Ramírez, et al. [16] have contributed to the understanding of Taylor series in solving definite integrals and evaluating functions. The current paper extends this knowledge by presenting additional examples, such as the Gaussian integral and sine integral, demonstrating the effectiveness of Taylor series in evaluating integrals lacking closed-form antiderivatives.

Asymptotic Behavior and Error Analysis

The concept of asymptotic behavior and error analysis in Taylor series is discussed by Mouze [14] and Georgiev and Erhan [6]. These works lay the groundwork for understanding the convergence criteria and error terms in Taylor series expansions. The present paper contributes by providing practical examples from physics and chemistry, showcasing how Taylor series aids in describing asymptotic behavior in real-world scenarios.

Applications in Fluid Dynamics

The paper references applications in fluid dynamics, aligning with works such as Estévez Schwarz and Lamour [5], who discuss explicit and implicit Taylor series methods for differential algebraic equations. However, the current paper extends beyond by introducing the application of Taylor series in describing the behavior of electric fields and molecular geometry optimization in chemistry.

Real Analysis and Poroelasticity

The exploration of Taylor series in real analysis, especially in the context of poroelasticity, connects with research by Bachi, et al. [2], who discuss statistical Taylor series expansion for uncertainty propagation. The current paper expands on this by highlighting the applications in petroleum geosciences, emphasizing the practicality of Taylor series in handling uncertainty in complex systems.

Computational Fluid Dynamics (CFD) and Taylor Weak Statement (TWS)

The paper touches upon computational fluid dynamics and the Taylor Weak Statement. While have discussed Taylor series solutions for various equations, the current paper extends this by applying the Taylor Weak Statement to analyze CFD outputs, demonstrating its robustness and accuracy.

In summary, this paper builds upon the foundational research in Taylor series applications, providing a comprehensive exploration of its diverse uses. It offers practical examples, extends existing concepts, and demonstrates the applicability of the Taylor series in solving complex problems across multiple scientific disciplines. The exploration of Taylor series applications aligns with and expands upon existing research in various scientific domains. Foundational works such as those by Veigend, et al. [20] and Bawazeer, et al. [3] provide insights into Taylor series-based numerical integration and radial basis function expansion. Guillot, et al. [7] and Ramírez, et al. [16] have contributed significantly to understanding Taylor series in

solving definite integrals and evaluating functions. The works by Mouze [14] and Georgiev and Erhan [6] lay the groundwork for error analysis and understanding the convergence criteria in Taylor series expansions. Estévez Schwarz and Lamour [5] discuss explicit and implicit Taylor series methods for differential algebraic equations, while Bachi, et al. [2] explore the statistical Taylor series expansion for uncertainty propagation. This paper builds upon these foundational studies, presenting practical examples and extending existing concepts to showcase the applicability of the Taylor series in solving complex problems across multiple scientific disciplines.

Taylor Series

The term Taylors series refers to an expansion of an ordinary function into a series of variable x , and then into a finite remainder term. Its coefficients are derivatives of the given function, and the subsequent terms involve the successive ones. The Taylor series is a finite set of numbers that were created by the expansion of Taylors. The magnitude of the remainder is the only concern. Given the interval between expansion a and b , the Lagrangian form is given as the remainder [6].

$$R_n = \frac{(x-a)^n}{n!} f^{(n)}(\xi)$$

a , is the reference point. The $f^{(n)}(\xi)$ is the n th derivative at a . When the expanding function is such that:

$$\lim_{n \rightarrow \infty} R_n = 0,$$

The Taylors series of the expanding functions becomes.

$$f(x) = \sum_{n=0}^{\infty} \frac{(x-a)^n}{n!} f^{(n)}(a).$$

Taylor series specifies the value of a function at one point, x . Setting the derivative operator, $D = d/dx$, the Taylor expansion becomes [11]:

$$f(x+h) = \sum_{n=0}^{\infty} \frac{h^n D^n}{n!} f(x) = e^{hD} f(x) [2]$$

Taylor series could also be written in the context of a complex variable [19]

Definite Integrals

Some functions do not have anti-derivatives. This makes it hard to evaluate their definite integrals. However, a series of representations can help solve this problem [7].

$$\int_0^1 \sin(x^2) dx$$

this integrand has no anti-derivative expressible in terms of familial functions. However, we know how to find its Taylor series:

$$\sin t = t - \frac{t^3}{3!} + \frac{t^5}{5!} - \frac{t^7}{7!} + \dots$$

if, we substitute $t = x^2$, then.

$$\sin(x^2) = x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} - \frac{x^{14}}{7!} + \dots$$

The Taylor series can then be integrated:

$$\int_0^1 \sin(x^2) dx = \frac{x^3}{3} - \frac{x^7}{7 \times 3!} + \frac{x^{11}}{11 \times 5!} - \frac{x^{15}}{15 \times 7!} + \dots$$

The series is composed of alternating terms and converges to a total of 0.31026.

The section on definite integrals introduces the challenge posed by functions lacking closed-form antiderivatives and highlights the utility of Taylor series in addressing this issue. To provide a more comprehensive understanding, let's delve into additional examples that demonstrate how the Taylor series can be effectively employed to evaluate integrals.

Example 1: The Gaussian Integral

Consider the Gaussian integral:

$$\int_{-\infty}^{\infty} e^{-x^2} dx$$

This integral is notorious for not having a closed-form antiderivative expressible in terms of elementary functions. However, we can use the Taylor series expansion for the exponential function to tackle this problem.

The Taylor series for e^{-x^2} is:

$$e^{-x^2} = 1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots$$

Now, substitute this series into the integral:

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \int_{-\infty}^{\infty} \left(1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots \right) dx$$

The integral of each term can be calculated individually, providing a series representation for the Gaussian integral. While this series converges slowly, it exemplifies how Taylor series can be a valuable tool for evaluating integrals without elementary antiderivatives.

Example 2: Sine Integral

Consider the integral:

$$\int_0^{\frac{\pi}{2}} \sin(x^2) dx$$

This integral does not possess a simple closed-form antiderivative. However, by expressing $\sin(x^2)$ using its Taylor series:

$$\sin(t) = t - \frac{t^3}{3!} + \frac{t^5}{5!} - \frac{t^7}{7!} + \dots$$

Substitute $t = x^2$ into the series and integrate each term individually. This yields a series representation for the sine integral, showcasing the practicality of Taylor series in evaluating integrals with challenging antiderivatives.

By incorporating these additional examples, the section on definite integrals

becomes more illustrative, demonstrating how Taylor series can be a powerful tool for tackling integrals with no closed-form antiderivatives.

Asymptotic Behavior

Error Analysis in Taylor Series

When utilizing a truncated Taylor series to approximate a function, it is essential to understand the associated error. The error term, often denoted as R_n , represents the difference between the actual function and its Taylor series approximation up to the n -th term.

$$R_n = \frac{(x-a)^{n+1}}{(n+1)!} f^{(n+1)}(\xi)$$

Here, a is the reference point, $f^{(n)}(\xi)$ is the n -th derivative at ξ , and ξ lies between a and x . The convergence of the Taylor series relies on the limit of this error term as n approaches infinity.

$$\lim_{n \rightarrow \infty} R_n = 0$$

Understanding this convergence criterion provides valuable insights into the accuracy of the Taylor series approximation.

Practical Examples

Let's explore practical examples from different scientific disciplines to illustrate the application of Taylor series in describing asymptotic behavior.

Physics - Electric Fields

Consider the electric field (E) obeying the inverse square law ($E = \frac{kq}{r^2}$). When the source is far away, an electric dipole moment ($E = \frac{kq}{(x-r)^2} + \frac{kq}{(x+r)^2}$) can be expanded using Taylor series. If we introduce xr into the series, we obtain an approximation for E :

$$E \approx \frac{4rq}{x^3}$$

This illustrates the use of Taylor series in describing the asymptotic behavior of electric fields as the distance from the source becomes significantly large.

Chemistry - Molecular Geometry Optimization

In chemistry, the equilibrium geometry of a molecule (U) can be approximated using Taylor series. Assuming initial coordinates x and y , the Taylor series approximation around (x_1, y_1) is expressed as:

$$U(x, y) \approx U(x_1, y_1) + \left. \frac{\partial U}{\partial x} \right|_{(x_1, y_1)} (x - x_1) + \left. \frac{\partial U}{\partial y} \right|_{(x_1, y_1)} (y - y_1) +$$

This application showcases how Taylor series assists in approximating molecular equilibrium positions.

The Taylor series can be used to describe the behavior of a given function in a sub domain. The electric field obey the inverse square law [13].

$$E = \frac{kq}{r^2}$$

The electric field is E, the charge is q, the distance from the source is r, and the proportionality is k. In this case, the electric field is far away from the source. A pair of charges placed side by side create an electric dipole moment [4].

$$E = \frac{kq}{(x-r)^2} + \frac{kq}{(x+r)^2}$$

An electric field is obtained from (10) after the terms in the denominator have been expanded.

$$E = \frac{kq}{x^2 \left(1 - \frac{r}{x}\right)^2} - \frac{kq}{x^2 \left(1 + \frac{r}{x}\right)^2}$$

The series Taylor can be expanded to add more numbers if xr is used.

$$\begin{aligned} \frac{kq}{\left(1 - \frac{r}{x}\right)^2} &= 1 + \frac{2r}{x} + \frac{3r^2}{x^2} + \frac{4r^3}{x^3} \\ \frac{kq}{\left(1 + \frac{r}{x}\right)^2} &= 1 - \frac{2r}{x} + \frac{3r^2}{x^2} - \frac{4r^3}{x^3} \end{aligned}$$

One Now Obtains

$$E \approx \frac{4rq}{x^3}$$

One of the most important factors that scientists consider when it comes to optimizing physical systems is the geometric design of their structures. In chemistry, for instance, a quasi-newton approach is used to calculate the equilibrium geometry of an atom [8]. For instance, let's assume that the molecule's geometry is U, and that its initial coordinates are the two x and y variables. The Taylor series is then used to approximate the molecule's equilibrium as shown next.

$$\begin{aligned} U(x, y) &\approx U(x_1, y_1) + \left. \frac{\partial U}{\partial x} \right|_{(x_1, y_1)} (x - x_1) \\ &+ \left. \frac{\partial U}{\partial y} \right|_{(x_1, y_1)} (y - y_1) + \frac{1}{2} \left. \frac{\partial^2 U}{\partial x^2} \right|_{(x_1, y_1)} (x - x_1)^2 \\ &+ \frac{1}{2} \left. \frac{\partial^2 U}{\partial y^2} \right|_{(x_1, y_1)} (y - y_1)^2 + \left. \frac{\partial^2 U}{\partial x \partial y} \right|_{(x_1, y_1)} (x - x_1)(y - y_1) \end{aligned}$$

The U(x) and y functions are well suited to the equilibrium position. The approximation of the quadratic function is also well within the minimum. If the U function is accurate, the Hessian matrix' elements will be constant in the region near the x_1, y_1 . One of the most challenging factors when it comes to calculating the second derivatives of a complex system is the time required to perform the procedure. This is because the ab initio method requires a lot of time to perform the optimization. For instance, if the U(x) form is written on the page, then the first approximation to its Hessian matrix element should be made [12, 15].

Application in Real Analysis

The properties of poroelasticity are useful in the design of various petroleum

geosciences applications. They are connected to the saturated elastic moduli of a porous rock by a Gassmann relation. Because of the uncertainty associated with the inputs, the error propagation in the estimates is usually an issue [2].

Monte-Carlo methods and Taylor's series expansion are two common error propagation techniques. The former requires a first-order analysis, while the latter usually only requires a computation. The formulas for analytical derivatives used to be incredibly complex, but they are now much easier to use with the help of modern symbolic packages. A numerical method for analyzing the effects of variable bathymetry on surface water waves is proposed. It can be used for both two- or three-dimensional flows as well as moving bottom topography. The method is based on a Hamiltonian system's boundary quantities [9].

The Dirichlet-Naumann operator is used in the analysis of boundary conditions such as crystal growth and fluid dynamics. It is computed using the Taylor series representation. The new stabilized forms of the Taylor terms are efficiently computed using a pseudo spectral method. The method is optimized for the order Taylor series expansion. The results show that the total computing error is 0.3 dB lower [18]. The proposed function generator has a maximum output range of more than 40 dB. Increasing the number of terms in the Taylor expansion can greatly reduce the total error. The expressions of the cumulants in terms of the Taylor series are based on the coefficients of the special function known as the Riemann Zeta. This allows for the calculation of an approximate form of the prime counting function [1, 18]. The method for obtaining the maximum number of cumulates is systematized, which yields the Taylor series expression for each of the various types of cumulants. The results are then analyzed by the Galerkin computational fluid dynamics algorithm. This method is ideal for analyzing the Incompressible Navier–Stokes (INS) equation. It is performed through the Taylor series expansion and a pseudo-limit process [3].

The Taylor Weak Statement is a spatially finite element analysis method that is used for analyzing the INS. It generates a CFD algorithm by implementing the Taylor Weak Statement. The output of this method is then analyzed by the generic node. The subsequent expansion of the Taylor series in wave number space affects the lower order error terms. Through the study of the complexity of the Taylor series, we have learned that it is an incredibly powerful tool for analyzing complex functions. Its applications in computational science and physical sciences are also widely acknowledged. The efficiency of the Taylor series in error determination, limit determination, and function optimization is evidenced by its numerous applications.

Function Approximation: Taylor series is extensively used for approximating functions. By truncating the series after a finite number of terms, a polynomial that closely resembles the original function within a certain interval is obtained. This is particularly useful in numerical methods for solving differential equations and performing numerical integration.

Solving Differential Equations: Many differential equations that model physical phenomena cannot be solved exactly. Taylor provides a method to find approximate solutions by expanding the unknown function as a series and solving the coefficients term by term. This approach is foundational in perturbation theory, where small deviations from known solutions are studied using series expansions.

Practical Uses in Physics and Engineering: In physics, Taylor series is used to approximate potential energy in systems around equilibrium points, leading to

linear approximations that simplify the analysis of physical systems. In engineering, it aids in the design and analysis of control systems, where small deviations from a set point are studied using linear approximations derived from Taylor series.

Computational Fluid Dynamics (CFD): Taylor series plays a crucial role in CFD, where it is used to develop numerical methods for solving fluid flow problems. The Taylor Weak Statement (TWS) method, for instance, is employed to analyze the Incompressible Navier-Stokes (INS) equation, providing robust and accurate solutions.

Error Analysis and Asymptotic Behavior

When utilizing a truncated Taylor series to approximate a function, understanding the associated error is essential. The error term R_n represents the difference between the actual function and its Taylor series approximation up to the n -th term:

$$R_n = \frac{(x - a)^{n+1}}{(n + 1)!} f^{(n+1)}(\xi)$$

where ξ lies between a and x . The convergence of the Taylor series relies on the limit of this error term as n approaches infinity. Practical examples from physics and chemistry will illustrate how Taylor series aids in describing asymptotic behavior in real-world scenarios.

Conclusion

The method for getting the maximum number of cations is systematized. It produces the Taylor series for each type of cation and then the k th one. The results of the Galerkin Fluid Dynamics algorithm are thoroughly analyzed by the Taylor series expansion followed by a pseudo-limit procedure. This is an ideal method for analyzing the INS equation due to its robust and high accuracy. The TWS method democratizes spatial elements in the implementation of INS. This allows the analysis of the results using a computational fluid dynamics algorithm. The amplification factor and phase velocity errors are then derived from the output of the algorithm for big and linear basis implementations. The changes in the Taylor series due to the wave space expansion affect the lower order error terms. Through its complexity, we have been able to show that the Taylor series is very effective in various applications. Its applications in computational science and physical sciences are also strong. The efficiency of the Taylor series in error determination, limit determination, and function optimization is shown to be outstanding. Taylor series is a fundamental tool in real analysis, providing powerful techniques for approximating functions, solving differential equations, and analyzing the convergence properties of functions. Its versatility and practicality make it an essential component of mathematical analysis, with applications across various scientific and engineering disciplines. Future research can explore advanced applications of Taylor series in modern computational methods, such as machine learning and data science.

Limitation

Though there are various applications and usage of numerical methods, these method's accuracy and precision are greatly impacted by certain limitations. The most

prominent issue is the truncation error linked with techniques of series expansion like the Taylor Series. Since approximations are numerical estimations that require a series to be truncated, the estimate may not result as expected, especially when working with rapid changes for functions or functions with singular points.

Another issue stems from intricacy of computation. Higher order Taylor series approximations, multi-variable functions and large-scale simulations incur hefty amounts of computation resources and time. This drawback is particularly apparent in applications that need results in real time since calculations have to be performed without delay. There are also problems regarding stability and convergence in numerical approximations. Even though Taylor series expansion tends to work well with smooth functions, it is not effective with all functions, thus, results cannot be dependent on all calculations. Discontinuous functions or functions with a small radius of convergence can cause the results to be off when numerical methods based on Taylor series are used.

Additionally, rounding metrics, which stem from limited precision calculations, are an issue for most numerical methods. Such errors often build up in iterative computations and pursue a deviation from the exact solution in the long iterative shifts which become much greater than they should be. The bottom line is, Taylor series method approximations and numerical methods are useful for solving complex mathematical problems, but their limitations require attention ensuring accurate and efficient computations. Everyone must deal with truncation errors, computational costs, convergence behavior, and rounding errors when using mathematical methods for numerical analysis in science and engineering.

References

- [1] A. Amarioarei, M. Paun, B. Strimbu, *Development of Nonlinear Parsimonious Forest Models Using Efficient Expansion of the Taylor Series: Applications to Site Productivity and Taper*, *Forests* **11** (2020), 458.
- [2] K. Bachi, K. Abbas, B. Heidergott, *Statistical Taylor series expansion: An approach for epistemic uncertainty propagation in Markov reliability models*, *RAIRO-Oper. Res.* **55** (2021), S593-S624.
- [3] S. A. Bawazeer, S. S. Baakeem, A. A. Mohamad, *New Approach for Radial Basis Function Based on Partition of Unity of Taylor Series Expansion with Respect to Shape Parameter*, *Algorithms (Basel)* **14** (2021), 1.
- [4] A. Elsaid, S. M. Helal, *Moving Taylor series for solving one-dimensional one-phase Stefan problem*, *Alexandria Engineering Journal* **61** (2022), 7121-7128.
- [5] D. Estévez Schwarz, R. Lamour, *Projected explicit and implicit Taylor series methods for DAEs*, *Numer. Algorithms* **88** (2021), 615-646.
- [6] S. G. Georgiev, I. M. Erhan, *The Taylor series method and trapezoidal rule on time scales*, *Appl. Math. Comput.* **378** (2020), 125200.
- [7] L. Guillot, B. Cochelin, C. Vergez, *A Taylor series-based continuation method for solutions of dynamical systems*, *Nonlinear Dyn.* **98** (2019), 2827-2845.
- [8] C.-H. He, Y. Shen, F.-Y. Ji, J.-H. He, *Taylor Series Solution for Fractal Bratu-type Equation Arising in Electrospinning Process*, *Fractals* **28** (2020), 2050011.
- [9] J.-H. He, *Taylor series solution for a third order boundary value problem arising in Architectural Engineering*, *Ain Shams Engineering Journal* **11** (2020), 1411-1414.
- [10] T. Insperger, *On the Approximation of Delayed Systems by Taylor Series Expansion*, *J. Comput. Nonlinear Dynam.* **10** (2015), 024503.

- [11] D. Jose, S. Sebastian, *Taylor series method in TDOA approach for indoor positioning system*, International Journal of Electrical and Computer Engineering (IJECE) **9** (2019), 3927-3933.
- [12] G. Jumarie, *Modified Riemann-Liouville derivative and fractional Taylor series of nondifferentiable functions further results*, Comput. Math. Appl. **51** (2006), 1367-1376.
- [13] G. Liu, J. Wang, *A relation spectrum inheriting Taylor series: muscle synergy and coupling for hand*, Frontiers of Information Technology & Electronic Engineering **23** (2022), 145-157.
- [14] A. Mouze, *Universal Taylor series with respect to a prescribed subsequence*, J. Math. Anal. Appl. **498** (2021), 124953.
- [15] L. Novák, D. Novák, *On Taylor Series Expansion for Statistical Moments of Functions of Correlated Random Variables*, Symmetry **12** (2020), 1379.
- [16] C. Ramírez, M. J. Rodríguez, B. D. Gomez, *Taylor series of Landauer conductance*, Physica E: Low-dimensional Systems and Nanostructures **124** (2020), 114213.
- [17] V. Satek, P. Veigend, G. Necasova, *Taylor Series Based Integration in Electric Circuits Simulations*, Advances in Electrical and Electronic Engineering **17** (2019), 352-359.
- [18] Q. Tang, J. Yang, X. He, W. Jia, Q. Zhang, H. Liu, *Nighttime image dehazing based on Retinex and dark channel prior using Taylor series expansion*, Comput. Vis. Image Underst. **202** (2021), 103086.
- [19] Y. Tong, S. Xiong, X. He, G. Pan, B. Zhu, *Symplectic neural networks in Taylor series form for Hamiltonian systems*, J. Comput. Phys. **437** (2021), 110325.
- [20] P. Veigend, G. Nečasová, V. Šátek, *Taylor Series Based Numerical Integration Method*, Open Computer Science **11** (2021), 60-69.
- [21] J. Yang, Y. Koutsawa, M. Potier-Ferry, H. Hu, *Changing variables in Taylor series with applications to PDEs*, Eng. Anal. Bound. Elem. **112** (2020), 77-86.
- [22] M. Yu, J. Lin, *Taylor series expansion scheme applied for solving population balance equation*, Reviews in Chemical Engineering **34** (2018), 561-594.